


# LABELLA



## Table variety

- ✓ Medium early yellow flesh variety
- ✓ High yielding
- ✓ Bright red skin
- ✓ Good dormancy
- ✓ PCN resistance (Ro1)
- ✓ Wart resistance (f1)



## Characteristics

### Physiology

Skin colour	Red
Flesh colour	Yellow
Tuber shape	Oval to long oval
Tuber shape uniformity	6
% 35/65	84%
Tubers per plant	14
Eye depth	8
Maturity	6,9
Dormancy	8
Storability	7
Bruising index (0-50)	5
Mechanical damage	7

### Yield - Quality

Relative yield	103
Dry matter content (UWW)	19,3% (351)
Cooking type	B
After cooking discoloration	6
Frying colour after harvest	6
Frying colour after storage	4

### Resistances

Foliage blight	2,8
Tuber blight	7
Common scab	6,4
Powdery scab	7
Globodera Ro 1 (A)	9
Wart f1 (D1)	10
Spraing (TRV)	8
Virus Y	6

## Growing advice (ware crop)

### Seed quantities and plant distance - Fertilizing

Size	kg/ha	75 cm ridge
28/35 mm	1.200	26-28
35/45 mm	2.300	30-32
45/50 mm	3.300	33-35
35/50 mm	2.600	31-33
50/60 mm cut	2.700	31-33

Always count the amount of tubers! Heavier soils require a closer planting distance.

Fertilizer	% compared to standard	Application 1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>th</sup>
N	110	3/4	1/4	
P <sub>2</sub> O <sub>5</sub>	100	4/4		
K <sub>2</sub> O	100	4/4		



- Can be grown on all soils.
- Planting on normal height in the ridge.
- Best results when planting in the white very small sprouts stage.
- Treatment against silverscab is advised.
- Prevent condensation before and during planting to avoid contamination.
- Does not seem sensible to Metribuzin.
- Weed control preferably before emergence.
- Start spraying against Phytophthora at 80% emergence.
- Early application of magnesium and manganese fertilizer has a positive effect.
- Second application of N-fertilizer during tuberization.
- Start spraying against Alternaria in time.
- Haulm killing after 110 days is advised.  
Lengthening the season by skipping haulm killing may cause a rough skin.
- Be careful with mother tubers and prevent damage during harvest.
- After harvest take care of good ventilation in storage until potatoes are dry.  
During long storage regular ventilation and gradually cooling down.

