

7 FOUR 7



Table variety

- ✓ Medium early white flesh variety
- ✓ Uniform long oval tubers
- ✓ High yielding
- ✓ Nice skin finish
- ✓ Suitable for washing
- ✓ PCN resistance (Ro1)



Characteristics

Physiology

Skin colour	Light yellow
Flesh colour	White
Tuber shape	Oval to long oval
Tuber shape uniformity	6
% 35/65	67%
Tubers per plant	13
Eye depth	8
Maturity	6,8
Dormancy	6
Storability	6
Bruising index (0-50)	2
Mechanical damage	7
Hollow hearts	9

Yield - Quality

Relative yield	113
Dry matter content (UWW)	17,1% (307)
Cooking type	AB
After cooking discoloration	7
Frying colour after harvest	4
Frying colour after storage	3

Resistances

Foliage blight	3,7
Tuber blight	7,7
Common scab	6,7
Powdery scab	7
Globodera Ro 1 (A)	9
Wart f1 (D1)	10
Spraing (TRV)	6,1
Virus Y	4



This information has been prepared to the best of our knowledge and belief and serves for orientation and decision-making. Since the potato is a natural product, we are not liable for any deviations. For questions or advice, please contact your cultivation supervisor or email info@denhartigh-potato.nl OCTOBER 2022 VERSION

Growing advice (ware crop)

Seed quantities and plant distance - Fertilizing

Size	kg/ha	75 cm ridge
28/35 mm	1.300	24-26
35/45 mm	2.300	28-30
45/50 mm	3.400	31-33
35/50 mm	2.700	29-31
50/60 mm cut	2.800	29-31

Always count the amount of tubers! Heavier soils require a closer planting distance.

Fertilizer	% compared to standard	Application		
		1 st	2 nd	3 th
N	100	2/4	1/4	1/4
P ₂ O ₅	100	4/4		
K ₂ O	110	4/4		



- Can be grown on all soil types.
- Best results when planting in the white very small sprouts stage.
- Treatment against silverscab is advised.
- Planting at normal height in de ridge.
- Do not use K-chloride as potassium source in spring.
- Does not seem susceptible to Metribuzin.
- Start spraying against Phytophthora at 80% emergence.
- Second application with N-fertilizer when plants start with tuberization.
- Start spraying against Alternaria in time.
- Third application with N fertilizer around flowering.
- Start spraying against tuber blight in time.
- After haulm killing, allow tuber skins to mature in the field before harvesting.
- After harvest take care of good ventilation in storage until potatoes are dry.